arXiv:1202.6265v1 [math-ph] 28 Feb 2012

Discretely holomorphic parafermions and integrable boundary conditions

Yacine Ikhlef

February 29, 2012

Abstract

In two-dimensional statistical models possessing a discretely holomorphic parafermion, we introduce a modified discrete Cauchy-Riemann equation on the boundary of the domain, and we show that the solution of this equation yields integrable boundary Boltzmann weights. This approach is applied to (i) the square-lattice O(n) loop model, where the exact locations of the special and ordinary transitions are recovered, and (ii) the Fateev-Zamolodchikov $\mathbb{Z}_N$ spin model, where a new rotation-invariant, integrable boundary condition is discovered for generic N.

1 Introduction

Discretely holomorphic parafermions (also called lattice parafermions) are lattice analogs for the holomorphic fields of Conformal Field Theory (CFT). They are lattice observables which satisfy a discrete version of the Cauchy-Riemann (CR) equation, and were recently identified in various two-dimensional critical lattice models, which are either spin models with a local interaction or loop models describing self-avoiding walks (SAWs) or extended interfaces. Generally, in spin models with Kramers-Wannier duality (see [1] for a recent review), a lattice parafermion is constructed [2, 3] by taking the product of lattice spin and disorder operators, providing an exact lattice analog for the operator product expansion (OPE) of the corresponding CFT [4]. In loop models, the lattice parafermion is expressed [2, 5] in terms of a single open path attached to the boundary of the system.

Lattice parafermions have found many applications to the rigorous study of lattice models (especially the Ising and SAW models) in the scaling limit, providing mathematical proofs for the exact results obtained by the Coulomb gas approach [6, 7] thirty years ago. In particular, the lattice parafermion is a crucial ingredient in the proofs [8, 9] of convergence of Ising domain walls to the Schramm-Loewner Evolution (SLE), and in the calculation of Ising correlation functions [10]. Also, for the SAW model on the honeycomb lattice, the lattice parafermion is essential to determine rigorously the location of the critical weights, both in the bulk [11] and on the boundary [12, 13].

This paper is concerned with lattice models at finite size, and addresses the relation between lattice parafermions and quantum integrability. Previous studies [2, 3, 5] have shown that many (spin or loop) statistical models admit a lattice parafermion exactly at their integrable point. In other words, the Boltzmann weights $\{W(\alpha)\}$ which satisfy the discrete CR equations on the rhombic lattice of angle α are also a solution of the Yang-Baxter (YB) equations, where α plays the role of the spectral parameter. In the present work, it is shown that a similar relation holds between the solutions of CR equations *on the boundary* and of Sklyanin's reflection equation [14] (or boundary YB equation).

Two distinct models are considered in this paper. First, for the O(n) loop model on the square lattice [15], following some ideas used in [12], local CR equations around boundary faces are introduced, and their solution is related to the boundary integrable weights found in [16]. When the rhombus angle is set to $\alpha = \pi/3$, one recovers the result of [12] for the honeycomb lattice. Then, we study the Fateev-Zamolodchikov $\mathbb{Z}_N$ spin model [17]. We introduce a similar boundary CR equation, and show that its solution yields new non-trivial integrable weights, associated to free BCs. In both cases, we use the lattice parafermions constructed in [3, 5] which satisfy bulk CR equations, and we define a new type of lattice domain Ω where some of the boundary faces have a differnet shape from the bulk ones.



Figure 1: The local configurations of the square-lattice O(n) model and the associated Boltzmann weights.



(a)

(b)

Figure 2: (a) A domain Ω_0 of the rhombic lattice with angle α. (b) A domain Ω of the rhombic lattice, where some boundary faces are replaced by triangles. In both cases, a is an arbitrary midedge of a boundary rhombus.

2 The O(n) model on the square lattice

2.1 Integrable weights and Cauchy-Riemann equation in the bulk

The O(n) model on the square lattice was introduced in [15]. It is a loop model, defined by the local configurations shown in Fig. 1, and the Boltzmann weight of a loop configuration G:

$$\Pi[G] = n^{N_\ell(G)} \times t^{N_t(G)} u_1^{N_{u_1}(G)} \dots w_2^{N_{w_2}(G)} ,$$

where $N_\ell(G)$ is the number of closed loops in G, and N_x is the number of faces with configuration x. The partition function is denoted by Z. If the loop fugacity is set to

$$n = -2\cos 4\lambda ,$$

the integrable weights are given by

$$\begin{array}{rcl} t & = & \sin(3\lambda - u)\sin u + \sin 2\lambda \sin 3\lambda \\ u_1 & = & -\sin(3\lambda - u)\sin 2\lambda \\ u_2 & = & -\sin u \sin 2\lambda \\ v & = & \sin(3\lambda - u)\sin u \\ w_1 & = & \sin(3\lambda - u)\sin(2\lambda - u) \\ w_2 & = & -\sin(\lambda - u)\sin u , \end{array} \tag{1}$$

where u is the spectral parameter. We first consider a domain Ω_0 of the rhombic lattice $\mathcal{R}_\alpha$ (see Fig. 2a), and recall the results on the lattice parafermion for the O(n) model. For a fixed edge a on the boundary, the parafermionic observable F_s is defined on the midedges $\{z\}$ of the rhombi as [5]

$$F_s(z) := \frac{1}{Z} \sum_{G \in \Gamma(a \to z)} \Pi[G] \; e^{-is\theta(G)} , \tag{2}$$

where $\Gamma(a \to z)$ is the set of loop configurations comprising an open path between a and z, $\theta(G)$ is the winding angle of the path included in G, and s is called the conformal spin. For a polygon P



Figure 3: The two possible configurations of a triangular face contributing to (8) for a fixed outer configuration.

with vertices given by the complex numbers $(p_1, \dots, p_m)$ ordered counter-clockwise, we define the discrete contour integral of a function f around P as

$$\sum_P f(z)\delta z := \sum_{j=1}^{m} (p_{j+1} - p_j) f\left(\frac{p_{j+1} + p_j}{2}\right) , \tag{3}$$

where $p_{m+1} := p_1$. We define the discrete CR equations as

$$\sum_{\diamond} F_s(z)\delta z = 0 , \tag{4}$$

where the left-hand side stands for the discrete integral around any given rhombus of Ω_0. Fixing the outer loop configuration, this gives the linear system:

$$\begin{array}{rcl} t + \mu u_1 - \mu\tau^{-1}u_2 - v &=& 0 \\ -\tau^{-1}u_1 + nu_2 + \tau\mu v - \mu\tau^{-1}(w_1 + nw_2) &=& 0 \\ nu_1 - \tau u_2 - \mu\tau^{-2}v + \mu(nw_1 + w_2) &=& 0 \\ -\mu\tau^{-2}u_1 + \mu\tau u_2 + nv - \tau^{-2}w_1 - \tau^2 w_2 &=& 0 , \end{array} \tag{5}$$

where we have set

$$\tau := e^{i\pi s} , \qquad \mu := e^{i(s+1)\alpha} .$$

In [5] it was shown that the system (5) has a nonzero solution iff $\cos 4\pi s = \cos 12\lambda$, and one can choose the value

$$s = \frac{3\lambda}{\pi} - 1 . \tag{6}$$

Moreover, the solution of (5) for $v \neq 0$ is given by the integrable weights (1), if one sets the spectral parameter to

$$u = (s+1)\alpha = \frac{3\lambda}{\pi}\,\alpha . \tag{7}$$

2.2 Boundary Cauchy-Riemann equation

We now consider a domain Ω where the interior faces are rhombi with a lower angle α, and the boundary faces are either rhombi or triangles (see Fig. 2b). Moreover, we assume that the initial orientation of the open path at the boundary point a is horizontal. The Boltzmann weights on a triangular face are y, r (see Fig. 1).

Let us introduce the boundary CR equation:

$$\mathrm{Re}\left[\sum_{\triangleleft} F_s(z)\delta z\right] = 0 , \tag{8}$$

where the discrete contour integral is around a triangular boundary face T. The configurations G in (2) which contribute to (8) are those where the open path ends on an edge of the boundary face T. Like in the bulk, one can fix the loop configuration outside T and sum over the internal configurations (see Fig. 3). This yields

$$\mathrm{Re}\left[-e^{\frac{i}{2}[(s+1)\alpha+\pi(1-s)]}r - e^{\frac{i}{2}[-(s+1)\alpha+\pi(1-s)]}y\right] = 0 . \tag{9}$$

We now have to choose an appropriate parameterisation for the spectral parameter. For this, we consider a particular instance of a domain Ω: the infinite strip in a diagonal direction of $\mathcal{R}_\alpha$. In [16], it is shown how to construct an integrable model on a strip with arbitrary, homogeneous



Figure 4: Left: the original square lattice $\mathcal{L}$ for the $\mathbb{Z}_N$ spins σ_j (full dots, full lines) and its dual lattice (empty dots, dotted lines). Right: the covering lattice $\mathcal{R}$.

spectral parameter in the bulk. If we denote $\check{R}$ and K the solutions to the bulk and boundary YB equations, the resulting model has bulk weights given by $\check{R}(u)$, and boundary weights given by $K(u/2)$. This suggests that in (9), one should set

$$u = \frac{1}{2}(s+1)\alpha$$

instead of (7). In contrast, the conformal spin belongs to the definition of the observable F_s, and we keep its value (6). The solution of (9) then reads

$$\begin{aligned} r(u) &= -\cos \tfrac{1}{2}(3\lambda + 2u) \\ y(u) &= \cos \tfrac{1}{2}(3\lambda - 2u) \, . \end{aligned} \tag{10}$$

These are exactly the boundary weights identifed in [16], which satisfy the boundary YB equation

$$\check{R}_{12}(u-v)K_2(u)\check{R}_{12}(u+v)K_2(v) = K_2(v)\check{R}_{12}(u+v)K_2(u)\check{R}_{12}(u-v)\, .$$

In [16], it is argued that (10) corresponds to the special surface transition in a certain regime of λ. Under the change $\lambda \to \lambda + \pi$, the bulk weights are invariant (up to irrelevant sign factors), whereas the boundary weights correspond to the ordinary surface transition.

Note that, if we specialise to $u = \lambda$ ($\alpha = \frac{\pi}{3}$), we recover the critical bulk and boundary weights of the O(n) model on the honeycomb lattice, which were found through a type of boundary CR equation in [12] for a specific domain Ω.

3 The Fateev-Zamolodchikov $\mathbb{Z}_N$ model

3.1 Definitions

The $\mathbb{Z}_N$ model consists of spins σ_j living on the sites of the square lattice $\mathcal{L}$ (full dots in Fig. 4), and taking the values:

$$\sigma_j \in \{1, \omega, \dots, \omega^{N-1}\}\, , \qquad \text{where} \qquad \omega = \exp\left(\frac{2i\pi}{N}\right)\, . \tag{11}$$

We shall then write $\sigma_j = \omega^{r_j}$, with $r_j \in \mathbb{Z}$. We also introduce the crossing parameter

$$\lambda := \frac{\pi}{2N}\, .$$

The Boltzmann weight of a spin configuration is:

$$\Pi[\sigma] = \prod_{\langle ij \rangle} W(\sigma_i, \sigma_j)\, , \qquad \text{where} \qquad W(\sigma_i, \sigma_j) = \begin{cases} W(u|r_i - r_j) & \text{if } \langle ij \rangle \text{ is horizontal,} \\ W(\lambda - u|r_i - r_j) & \text{if } \langle ij \rangle \text{ is vertical,} \end{cases} \tag{12}$$

where u is the spectral parameter. Moreover, we restrict to a periodic and symmetric function $W(u|r)$:

$$W(u|r+N) = W(u|-r) = W(u|r)\, . \tag{13}$$

The partition function is then defined as

$$Z = \sum_{\{\sigma\}} \Pi[\sigma]\, , \tag{14}$$



Figure 5: The star-triangle relation as a special case of the Yang-Baxter equation.

and a typical correlation function of n spin operators is of the form

$$\langle \sigma_{j_1}^{k_1} \sigma_{j_2}^{k_2} \dots \sigma_{j_n}^{k_n} \rangle := \frac{1}{Z} \sum_{\{\sigma\}} \Pi[\sigma] \sigma_{j_1}^{m_1} \sigma_{j_2}^{m_2} \dots \sigma_{j_n}^{m_n} \,,$$

where $\{j_1, \dots, j_n\}$ are n sites of the lattice $\mathcal{L}$, and $\{m_1, \dots, m_n\}$ are n arbitrary integers. As a consequence of the symmetries (13) of the Boltzmann weights, the model enjoys Kramers-Wannier duality, and in particular one may define dual spin operators, also called disorder operators $\mu^{(m)}$. These live on the dual lattice (empty dots in Fig. 4), and are defined through their correlation functions. For the purposes of the present study, it is sufficient to define correlation functions involving two μ operators. Let k and ℓ be sites of the dual lattice $\mathcal{L}'$, and $\gamma_{k\ell}$ an oriented path from k to ℓ on $\mathcal{L}'$. One then defines:

$$\langle \mu_k^{(-m)} \mu_\ell^{(m)} (\dots) \rangle := \frac{1}{Z} \sum_{\{\sigma\}} \prod_{\langle ij \rangle \notin \gamma_{k\ell}^\perp} W(\sigma_i, \sigma_j) \prod_{\langle ij \rangle \in \gamma_{k\ell}^\perp} W(\sigma_i, \omega^m \sigma_j) \, (\dots) \,, \tag{15}$$

where the dots denote any function of the spins, $\gamma_{k\ell}^\perp$ is the set of edges of $\mathcal{L}$ which cross $\gamma_{k\ell}$, and we have taken the convention that σ_i (σ_j) is on the left (right) of $\gamma_{k\ell}$. The symmetries (13) imply that $\langle \mu_k^{(-m)} \mu_\ell^{(m)} \rangle$ is independent of the choice of the path $\gamma_{k\ell}$.

In the following, we shall consider a deformation of the square lattice $\mathcal{L}$ into a rectangular lattice of aspect ratio $\tan \frac{\alpha}{2}$. The union of $\mathcal{L}$ and its dual is then a rhombic lattice of angle α, called the covering lattice and denoted $\mathcal{R}_\alpha$.

Later in this Section, we will use the discrete Fourier transform

$$\widehat{f}(k) := \sum_{r=0}^{N-1} \omega^{kr} f(r) \,, \qquad f(r) := \frac{1}{N} \sum_{r=0}^{N-1} \omega^{-kr} \widehat{f}(k) \,. \tag{16}$$

3.2 Integrable weights and Cauchy-Riemann equation in the bulk

The integrability condition for a periodic system (*i.e.* the commutation of transfer matrices on the cylinder) has the form of a star-triangle equation

$$\begin{aligned} &\sum_{r'=0}^{N-1} W(\lambda - u | r_1 - r') W(u + v | r_2 - r') W(\lambda - v | r_3 - r') \\ &\qquad = W(u | r_2 - r_3) W(\lambda - u - v | r_1 - r_3) W(v | r_1 - r_2) \,, \end{aligned} \tag{17}$$

which can be formulated as the Yang-Baxter equation on the covering lattice (see Fig. 5), and a solution to this equation was given in [17]:

$$W(u|0) = 1 \,, \qquad W(u|r) = \prod_{p=0}^{r-1} \frac{\sin[(2p+1)\lambda - u]}{\sin[(2p+1)\lambda + u]} \quad \text{for } r \in \{1, \dots, N-1\}. \tag{18}$$

This solution is self-dual, in the sense that

$$\widehat{W}(u|k) = W(\lambda - u|k) \,. \tag{19}$$

The construction [3] of the discrete parafermion in the $\mathbb{Z}_N$ model is the lattice version of the parafermionic current in the $\mathbb{Z}_N$ CFT [4]. In the lattice model, as well as in the CFT, the



Figure 6: A face of the rhombic covering lattice $\mathcal{R}_\alpha$ in the $\mathbb{Z}_N$ model. The dotted line represents an arbitray path γ on $\mathcal{L}'$ from $\mu(a)$ to μ_1.

parafermion ψ is obtained as the product of a spin operator σ and a disorder operator μ on adjacent sites. Consider a point z on the middle of an edge of $\mathcal{R}_\alpha$, and denote $j(z), k(z)$ the sites of $\mathcal{L}, \mathcal{L}'$ adjacent to z. We shall use the short-hand notations

$$\sigma(z) := \sigma_{j(z)}\,, \qquad \mu(z) := \mu^{(1)}_{k(z)}\,, \qquad \mu^*(z) := \mu^{(-1)}_{k(z)}\,.$$

The lattice parafermion $\psi(z)$ is defined as [3]

$$\psi(z) := \sigma(z)\ \mu(z)\,. \tag{20}$$

The observable $F_s(z)$ is then defined by fixing a point a on the boundary, and setting

$$\begin{aligned} F_s(z) &:= \langle e^{-is\theta(a,z)}\psi^*(a)\psi(z)\rangle \\ &= \langle e^{-is\theta(a,z)}\sigma^*(a)\sigma(z)\mu^*(a)\mu(z)\rangle\,, \end{aligned} \tag{21}$$

where $\theta(a,z)$ is the angle between the vectors $\overrightarrow{k(a)j(a)}$ and $\overrightarrow{k(z)j(z)}$, and s is the conformal spin. As in the O(n) model, one defines the discrete CR equation on $\mathcal{R}_\alpha$ as

$$\sum_{\diamond} F_s(z)\delta z = 0\,, \tag{22}$$

which can be written (see Fig. 6)

$$e^{-s\theta(a)}\ \langle\sigma^*(a)\sigma_1\mu^*(a)\mu_1\left[(e^{-iu} - e^{iu}\sigma_1^*\sigma_2) + (e^{iu+i\pi s} - e^{-iu-i\pi s}\sigma_1^*\sigma_2)\mu_1^*\mu_2\right]\rangle = 0\,, \tag{23}$$

where $\theta(a)$ is the angle between $\overrightarrow{k(a)j(a)}$ and the horizontal, and we have set

$$u = \frac{1}{2}(1-s)(\pi-\alpha)\,. \tag{24}$$

From the definition of μ, (23) reads

$$\begin{aligned} &\frac{1}{Z}\sum_{\{\sigma\}}\sigma^*(a)\sigma_1 \prod_{\langle ij\rangle\notin\gamma^\perp\cup\{\langle 12\rangle\}} W(\sigma_i,\sigma_j) \prod_{\langle ij\rangle\in\gamma^\perp} W(\sigma_i,\omega\sigma_j) \\ &\quad\times\left[(e^{-iu} - e^{iu}\sigma_1^*\sigma_2)W(\sigma_1,\sigma_2) + (e^{iu+i\pi s} - e^{-iu-i\pi s}\sigma_1^*\sigma_2)W(\sigma_1,\omega\sigma_2)\right] = 0\,, \end{aligned} \tag{25}$$

where γ is an arbitrary path on $\mathcal{L}'$ going from $\mu(a)$ to μ_1 (see Fig. 6). The bracket in (25) may be written as

$$I(r) := (e^{-iu} - e^{iu}\omega^r)W(u|r) + (e^{iu+i\pi s} - e^{-iu-i\pi s}\omega^r)W(u|r+1)\,, \tag{26}$$

where

$$\sigma_1^*\sigma_2 = \omega^r\,.$$

Therefore, a sufficient condition for (23) to hold is that $I(r)$ vanishes for all r, which yields the recursion relation

$$W(u|r+1) = W(u|r)\times(-e^{i\pi s+\frac{i\pi}{N}})\ \frac{\sin\left[\frac{(2r+1)\pi}{2N} - u\right]}{\sin\left[\frac{(2r+1)\pi}{2N} + u\right]}\,. \tag{27}$$



Figure 7: The two-row transfer matrix along one direction of the covering lattice, with open boundary conditions. Thick lines denote interaction terms between the spins.



Figure 8: The boundary Yang-Baxter equation for the $\mathbb{Z}_N$ model. Thick lines denote interaction between the spins, and spins joined by a double line must be equal.

Compatibility of (27) with the symmetries (13) fixes the value of the spin:

$$s = 1 - \frac{1}{N}\,, \tag{28}$$

and for this value, the solution of (27) is given by the integrable weights (18).

As explained in [3], each of the $(N-1)$ discrete parafermions $\psi^{(m)}(z) = \sigma^m(z)\mu^{(m)}(z)$ with charge $m \in \{1, \dots, N-1\}$ satisfies the CR equations, but for a different function $W^{(m)}$, and a conformal spin $s_m = m(N-m)/N$. In fact, $W^{(m)}$ is simply related to W (18) by the global transformation $\sigma_j \to \sigma_j^m$. Hence, one can restrict to the case $m = 1$ exposed above, without loss of generality.

3.3 Boundary Yang-Baxter equation

Following [18], we consider the boundary conditions which ensure the commutation of two-row transfer matrices (see Fig. 7). We denote $Y_{R,L}(\sigma_i, \sigma_j)$ the Boltzmann weight on the right and left boundaries, and the partition function on the strip reads

$$Z_{\text{strip}} = \sum_{\{\sigma\}} \prod_{\langle ij\rangle\in\text{bulk}} W(\sigma_i, \sigma_j) \prod_{\langle ij\rangle\in\text{right bound.}} Y_R(\sigma_i, \sigma_j) \prod_{\langle ij\rangle\in\text{left bound.}} Y_L(\sigma_i, \sigma_j)\,. \tag{29}$$

In [18], fixed-spin integrable boundary conditions (*i.e.* $Y(\sigma_i, \sigma_j) = \delta_{r_i,\alpha}\delta_{r_j,\beta}$ with α, β fixed) were considered, and the associated surface free energies were obtained. In the present work, we address the boundary YB equation for rotation-invariant boundary conditions, *i.e.* an interaction of the form

$$Y(\sigma_i, \sigma_j) = Y(u, \xi | r_i - r_j)\,, \tag{30}$$

where ξ is a complex parameter associated to each boundary (*boundary field*), and Y satisfies the symmetries (13). The boundary YB equation then reads (see Fig. 8)

$$\begin{aligned} &W(u-v|r_1-r_2) \sum_{r'=0}^{N-1} Y(u,\xi|r_1-r')W(u+v|r_2-r')Y(v,\xi|r_3-r') \\ &= W(u-v|r_2-r_3) \sum_{r'=0}^{N-1} Y(v|r_1-r')W(u+v|r_2-r')Y(u,\xi|r_3-r')\,, \end{aligned} \tag{31}$$



Figure 9: A boundary face in the $\mathbb{Z}_N$ model. Like in Fig. 6, the dotted line represents an arbitray path γ on $\mathcal{L}'$ from $\mu(a)$ to μ_1. In the discrete contour integral (33), only the edges represented by single lines contribute.

or, in Fourier space:

$$\widehat{Y}(v,\xi|k)\sum_{m=0}^{N-1}\widehat{W}(u-v|m)\widehat{Y}(u,\xi|\ell-m)\widehat{W}(u+v|k+\ell-m)$$

$$=\widehat{Y}(v,\xi|\ell)\sum_{m=0}^{N-1}\widehat{W}(u-v|m)\widehat{W}(u+v|k+\ell-m)\widehat{Y}(u,\xi|k-m)\,. \qquad (32)$$

These functional equations for Y are difficult to solve for general N, but we shall now show that a nontrivial solution can be found by use of the boundary CR equation.

3.4 Boundary Cauchy-Riemann equation

We consider boundary faces in the shape of a pentagon, determined by the angles α and β (see Fig. 9). The contour integral of F_s around such a face P reads:

$$\sum_P F_s(z)\delta z = e^{-s\theta(a)}\ \langle \psi^*(a)\sigma_1\mu_1\left[(e^{-2iu}-e^{2iu}\sigma_1^*\sigma_2)+(e^{i\xi+i\pi s}-e^{-i\xi-i\pi s}\sigma_1^*\sigma_2)\mu_1^*\mu_2\right]\rangle\,, \qquad (33)$$

where we have set[1]

$$u=\frac{1}{4}(1-s)(\pi-\alpha)\,,\qquad \text{and}\qquad \xi=\frac{1}{2}(1-s)(\pi-\beta)\,. \qquad (34)$$

Like for the bulk CR equation, using the notation $\sigma_1^*\sigma_2 := \omega^r$, (33) is rewritten as

$$\sum_P F_s(z)\delta z = \frac{e^{-s\theta(a)}}{Z}\sum_{\{\sigma\}}\sigma^*(a)\sigma_1\prod_{\langle ij\rangle\notin\gamma^\perp}W(\sigma_i,\sigma_j)\prod_{\langle ij\rangle\in\gamma^\perp}W(\sigma_i,\omega\sigma_j)\times J(r)\,,$$

where $J(r)$ is defined as

$$J(r) := (e^{-2iu}-e^{2iu}\omega^r)Y(u,\xi|r)+(e^{i\xi+i\pi s}-e^{-i\xi-i\pi s}\omega^r)Y(u,\xi|r+1)\,, \qquad (35)$$

and its Fourier transform is

$$\widehat{J}(k) := (e^{-2iu}+e^{i\xi+i\pi s}\omega^{-k})\widehat{Y}(u,\xi|k)-(e^{2iu}+e^{-i\xi-i\pi s}\omega^{-k-1})\widehat{Y}(u,\xi|k+1)\,. \qquad (36)$$

In analogy with the O(n) model, we introduce a boundary CR equation of the form

$$\mathrm{Re}\left[e^{i\varphi(k)}\widehat{J}(k)\right]=0\,, \qquad (37)$$

where $\varphi(k)$ is real. It turns out that Y only satisfies the symmetry relations (13) for the choice $\varphi(k)=(4k+2)\lambda$, and the solution of (37) then reads

$$\widehat{Y}(u,\xi|k)=\begin{cases}1 & \text{for } k=0.\\ \displaystyle\prod_{\ell=0}^{k-1}\frac{\sin\left[(2\ell+1)\lambda-u+\xi\right]\sin\left[(2\ell+1)\lambda-u-\xi\right]}{\sin\left[(2\ell+1)\lambda+u+\xi\right]\sin\left[(2\ell+1)\lambda+u-\xi\right]} & \text{for } k\in\{1,\dots,N-1\}.\end{cases} \qquad (38)$$

A remarkable fact is that *the weights* (38) *are also a solution of the boundary YB equation* (32). The solutions $W^{(m)}, Y^{(m)}$ associated to the other parafermions $\psi^{(m)}$ also satisfy the boundary YB equation, and they are related to (18)–(38) by a simple transformation $\sigma_j \to \sigma_j^m$.

---

[1] Like for the O(n) model, for a given angle α, the spectral parameter for boundary weights is half the bulk value (24).

3.5 Physical interpretation of the integrable boundary condition

To indentify the BC corresponding to (38), we consider the *diagonal-to-diagonal* transfer matrix, which is constructed from the two-row transfer matrix by the procedure described in [16]. If the two-row transfer matrix with horizontal spectral parameter u, vertical spectral parameters $(v_1, \dots, v_N)$ and boundary parameters $\xi_{L,R}$ is denoted

$$t(u, \xi_L, \xi_R | v_1, \dots, v_N) \,,$$

then the diagonal-to-diagonal transfer matrix is given by

$$t_d(u, \xi_L, \xi_R) := t\left(\frac{u}{2}, \xi_L, \xi_R \middle| \, \frac{u}{2}, -\frac{u}{2}, \dots, \frac{u}{2}, -\frac{u}{2}\right) \,.$$

In this setting, the bulk weights are given by $W(u|\cdot)$ for horizontal edges and $W(\lambda - u|\cdot)$ for vertical edges, whereas the left and right boundary weights read

$$Y(\lambda - u/2, \xi_L|r) W(\lambda - u|r) \,, \qquad \text{and} \qquad Y(u/2, \xi_R|r) \,.$$

The latter coincide for the value $\xi_L = \xi_R = u/2$, where both are proportional to $W(\lambda - u|r)$. Hence, for this choice of boundary parameters ξ_L, ξ_R, the boundary weights are equal to the bulk weights in the vertical direction, and thus (38) corresponds to free boundary conditions for the spins $\{\sigma_j\}$.

4 Conclusions

On the two examples we have studied, we have shown that the integrable boundary weights can be obtained by imposing a simple linear condition – which we have called the boundary CR equation – on the discrete contour integral

$$\sum_P F_s(z) \delta z \,,$$

where P stands for a modified boundary face. In the case of the O(n) model, we have recovered the integrable boundary weights which were obtained [16] through the mapping to the 19-vertex model, whereas for the $\mathbb{Z}_N$ model, we have found new integrable boundary weights. However, unlike the bulk CR equations, where the conformal spin s is generally obtained by simple consistency conditions, it is not clear yet how to extract conformal data from the boundary CR equations.

Our approach is quite general, and is likely to extend to any lattice model where a solution of the bulk CR equation has been identified. Also, an interesting continuation of this work would be to study the boundary CR equations in loop models with "non-trivial" integrable boundary conditions, *i.e.* where loops touching the boundary get a different weight [19].

Finally, an important possible application of our results would be to set up proofs of convergence to variants of SLE as in [8, 9], in the presence of various integrable boundary conditions.

Acknowledgements. The author thanks J.L. Jacobsen and M.A. Rajabpour for stimulating discussions, and a careful reading of the manuscript. This work was supported by the European Research Council (grant CONFRA 228046).

References

[1] J. Dubédat, ArXiv e-prints (2011), 1112.4404.

[2] V. Riva and J. Cardy, J. Stat. Mech. **2006**, P12001 (2006).

[3] M. A. Rajabpour and J. Cardy, J. Phys. A **40** (2007), 14703.

[4] A. B. Zamolodchikov and V. A. Fateev, Sov. Phys. JETP **62**, 215 (1985).

[5] Y. Ikhlef and J. Cardy, J. Phys. A **42** (2009), 102001.

[6] B. Nienhuis, J. Stat. Phys. **34**, 731 (1984).

[7] P. Di Francesco, H. Saleur, and J.-B. Zuber, J. Stat. Phys. **49**, 57 (1987).

[8] D. Chelkak and S. Smirnov, ArXiv e-prints (2009), 0910.2045.

[9] C. Hongler and K. Kytölä, ArXiv e-prints (2011), 1108.0643.

[10] D. Chelkak, C. Hongler, and K. Izyurov, ArXiv e-prints (2012), 1202.2838.

[11] H. Duminil-Copin and S. Smirnov, ArXiv e-prints (2010), 1007.0575.

[12] N. R. Beaton, J. de Gier, and A. J. Guttmann, ArXiv e-prints (2011), 1109.0358.

[13] N. R. Beaton, A. J. Guttmann, and I. Jensen, J. Phys. A **45**, 055208 (2012).

[14] E. K. Sklyanin, J. Phys. A **21**, 2375 (1988).

[15] B. Nienhuis, Physica A **163**, 152 (1990).

[16] C. Yung and M. Batchelor, Nucl. Phys. B **435**, 430 (1995).

[17] V. Fateev and A. Zamolodchikov, Phys. Lett. A **92**, 37 (1982).

[18] Y.-K. Zhou, Nucl. Phys. B **487**, 779 (1997).

[19] J. Dubail, J. L. Jacobsen, and H. Saleur, Phys. Rev. Lett. **103** (2009), 145701.